

The Advisory Committee on Immunization Practices' Interim Recommendation for Use of Pfizer-BioNTech COVID-19 Vaccine — United States, December 2020

Sara E. Oliver, MD¹; Julia W. Gargano, PhD¹; Mona Marin, MD¹; Megan Wallace, DrPH^{1,2}; Kathryn G. Curran, PhD¹; Mary Chamberland, MD^{1,3}; Nancy McClung, PhD¹; Doug Campos-Outcalt, MD⁴; Rebecca L. Morgan, PhD⁵; Sarah Mbaeyi, MD¹; José R. Romero, MD⁶; H. Keipp Talbot, MD⁷; Grace M. Lee, MD⁸; Beth P. Bell, MD⁹; Kathleen Dooling, MD¹

On December 13, 2020, this report was posted as an MMWR Early Release on the MMWR website (<https://www.cdc.gov/mmwr>).

On December 11, 2020, the Food and Drug Administration (FDA) issued an Emergency Use Authorization (EUA) for the Pfizer-BioNTech COVID-19 (BNT162b2) vaccine (Pfizer, Inc; Philadelphia, Pennsylvania), a lipid nanoparticle-formulated, nucleoside-modified mRNA vaccine encoding the prefusion spike glycoprotein of SARS-CoV-2, the virus that causes coronavirus disease 2019 (COVID-19) (1). Vaccination with the Pfizer-BioNTech COVID-19 vaccine consists of 2 doses (30 µg, 0.3 mL each) administered intramuscularly, 3 weeks apart. On December 12, 2020, the Advisory Committee on Immunization Practices (ACIP) issued an interim recommendation* for use of the Pfizer-BioNTech COVID-19 vaccine in persons aged ≥16 years for the prevention of COVID-19. To guide its deliberations regarding the vaccine, ACIP employed the Evidence to Recommendation (EtR) Framework,[†] using the Grading of Recommendations, Assessment, Development and Evaluation (GRADE) approach.[§] The recommendation for the Pfizer-BioNTech COVID-19 vaccine should be implemented in conjunction with ACIP's interim recommendation for allocating initial supplies of COVID-19 vaccines (2). The ACIP recommendation for the use of the Pfizer-BioNTech COVID-19 vaccine under EUA is interim and will be updated as additional information becomes available.

Since June 2020, ACIP has convened nine public meetings to review data on the epidemiology of COVID-19 and the potential use of COVID-19 vaccines, including the Pfizer-BioNTech COVID-19 vaccine (3). Within the EtR Framework, ACIP considered the importance of the public health problem of COVID-19, as well as issues of resource use, benefits and harms, patients' values and preferences, acceptability, feasibility, and equity for the Pfizer-BioNTech COVID-19 vaccine. To inform the EtR Framework, the COVID-19 Vaccines Work Group, comprising experts in infectious disease, vaccinology, vaccine safety, public health, and ethics, held 27 meetings

to review COVID-19 surveillance data, evidence for vaccine efficacy and safety, and implementation considerations for COVID-19 vaccines, including the Pfizer-BioNTech COVID-19 vaccine. After a systematic review of the literature, the Work Group used the GRADE approach to assess the certainty of evidence for outcomes related to the vaccine, rated on a scale of 1 (high certainty) to 4 (very low certainty) (4). Work Group conclusions regarding the evidence for the Pfizer-BioNTech COVID-19 vaccine were presented to ACIP at public meetings.

The body of evidence for the Pfizer-BioNTech COVID-19 vaccine was primarily informed by one large, randomized, double-blind, placebo-controlled Phase II/III clinical trial that enrolled >43,000 participants (median age = 52 years, range = 16–91 years) (5,6). Interim findings from this clinical trial, using data from participants with a median of 2 months of follow-up, indicate that the Pfizer-BioNTech COVID-19 vaccine was 95.0% effective (95% confidence interval = 90.3%–97.6%) in preventing symptomatic laboratory-confirmed COVID-19 in persons without evidence of previous SARS-CoV-2 infection. Consistent high efficacy (≥92%) was observed across age, sex, race, and ethnicity categories and among persons with underlying medical conditions as well as among participants with evidence of previous SARS-CoV-2 infection. Although numbers of observed hospitalizations and deaths were low, the available data were consistent with reduced risk for these severe outcomes among vaccinated persons compared with that among placebo recipients. Among vaccine recipients, reactogenicity symptoms, defined as solicited local injection site or systemic reactions during the 7 days after vaccination, were frequent and mostly mild to moderate. Systemic adverse reactions were more commonly reported after the second dose than after the first dose and were generally more frequent and severe in persons aged 18–55 years than in those aged >55 years. Systemic adverse reactions had a median onset of 1–2 days after vaccine receipt and resolved in a median of 1 day. Severe local and systemic adverse reactions (grade ≥3, defined as interfering with daily activity) occurred more commonly in vaccine recipients than in placebo recipients. Among vaccine recipients, 8.8% reported any grade ≥3 reaction; the most common symptoms were fatigue (4.2%), headache (2.4%), muscle pain (1.8%), chills (1.7%), and injection site pain

* On December 12, 2020, ACIP voted 11–0 (three recusals) in favor of the interim recommendation for use of Pfizer-BioNTech COVID-19 vaccine. Three ACIP members recused themselves because of participation in clinical trials and/or other studies involving companies producing COVID-19 vaccines.

[†] <https://www.cdc.gov/vaccines/acip/recs/grade/downloads/ACIP-evidence-rec-frame-508.pdf>.

[§] <https://www.cdc.gov/vaccines/acip/recs/grade/about-grade.html>.

(1.4%). Generally, grade ≥ 3 reactions were more commonly reported after the second dose than after the first dose and were less prevalent in older than in younger participants. Serious adverse events[‡] were observed in a similar proportion of vaccine (0.6%) and placebo (0.5%) recipients and encompassed medical events occurring at a frequency similar to that within the general population (6). No specific safety concerns were identified in subgroup analyses by age, race, ethnicity, underlying medical conditions, or previous SARS-CoV-2 infection. A detailed summary of safety data, including information on reactogenicity, is available at <https://www.cdc.gov/vaccines/covid-19/info-by-manufacturer/pfizer/reactogenicity.html>.

From the GRADE evidence assessment, the level of certainty for the benefits of the Pfizer-BioNTech COVID-19 vaccine was type 1 (high certainty) for the prevention of symptomatic COVID-19. Evidence was type 3 (low certainty) for the estimate of prevention of COVID-19–associated hospitalization and type 4 (very low certainty) for the estimate of prevention of death. Data on hospitalizations and deaths are limited at this time, but a vaccine that effectively prevents symptomatic infection is expected to also prevent hospitalizations and deaths. Regarding potential harms after vaccination, evidence was type 2 (moderate certainty) for serious adverse events and type 1 (high certainty) for reactogenicity. No data were available to assess the efficacy for prevention of asymptomatic SARS-CoV-2 infection. Data reviewed within the EtR Framework supported the use of the Pfizer-BioNTech COVID-19 vaccine. ACIP determined that COVID-19 is a major public health problem and that use of the Pfizer-BioNTech COVID-19 vaccine is a reasonable and efficient allocation of resources. Whereas there might be uncertainty in how all populations value the vaccine, it was determined that for most populations, the desirable effects outweigh the undesirable effects. The vaccine is probably acceptable to implementation stakeholders and feasible to implement in spite of difficult ultracold-chain storage and requirements for handling and administration. These requirements could limit the availability of the Pfizer-BioNTech COVID-19 vaccine to some populations thereby negatively impacting health equity. Therefore, efforts should be made to overcome these challenges and advance health equity. The GRADE evidence profile and EtR supporting evidence are available at <https://www.cdc.gov/vaccines/acip/recs/grade/covid-19-pfizer-biontech-vaccine.html> and <https://www.cdc.gov/vaccines/acip/recs/grade/covid-19-pfizer-biontech-etr.html>.

[‡]Serious adverse events are defined as any untoward medical occurrence that results in death, is life-threatening, requires inpatient hospitalization or prolongation of existing hospitalization, or results in persistent disability/incapacity.

Before vaccination, the EUA Fact Sheet should be provided to recipients and caregivers. Providers should counsel Pfizer-BioNTech COVID-19 vaccine recipients about expected systemic and local reactogenicity. Additional clinical considerations, including details of administration and use in special populations (e.g., persons who are pregnant or immunocompromised or who have severe allergies) are available at <https://www.cdc.gov/vaccines/covid-19/info-by-manufacturer/pfizer/clinical-considerations.html>. Additional studies of safety and effectiveness are planned after authorization and will be important to inform future ACIP recommendations as well as increase public confidence in the COVID-19 vaccination program. The interim recommendation and clinical considerations are based on use of the Pfizer-BioNTech COVID-19 vaccine under an EUA and might change as more evidence becomes available. ACIP will continue to review additional data as they become available; updates to recommendations or clinical considerations will be posted on the ACIP website (<https://www.cdc.gov/vaccines/hcp/acip-recs/vacc-specific/covid-19.html>).

Reporting of Vaccine Adverse Events

Adverse events that occur in a recipient after receipt of COVID-19 vaccine should be reported to the Vaccine Adverse Events Reporting System (VAERS). FDA requires that vaccination providers report vaccination administration errors, serious adverse events, cases of multisystem inflammatory syndrome, and cases of COVID-19 that result in hospitalization or death after administration of COVID-19 vaccine under EUA. Reporting is encouraged for any clinically significant adverse event, whether or not it is clear that a vaccine caused the adverse event. Information on how to submit a report to VAERS is available at <https://vaers.hhs.gov/index.html> or 1-800-822-7967. In addition, CDC has developed a new, voluntary smartphone-based tool, v-safe, that uses text messaging and web surveys to provide near real-time health check-ins after patients receive COVID-19 vaccination. The CDC/v-safe call center follows up on reports to v-safe that indicate a medically significant health impact to collect additional information for completion of a VAERS report. Information on v-safe is available at <https://www.cdc.gov/vsafe>.

Acknowledgments

Voting members of the Advisory Committee on Immunization Practices: Robert L. Atmar, Baylor College of Medicine; Kevin A. Ault, University of Kansas Medical Center; Lynn Bahta, Minnesota Department of Health; Henry Bernstein, Zucker School of Medicine at Hofstra/Northwell Cohen Children's Medical Center; Sharon E. Frey, Saint Louis University Medical School; Paul Hunter, City of Milwaukee Health Department; Veronica V. McNally, Franny Strong Foundation; Katherine A. Poehling, Wake Forest School of Medicine;

Pablo J. Sánchez, The Research Institute at Nationwide Children's Hospital; Peter Szilagyi, University of California, Los Angeles. Members of the Advisory Committee on Immunization Practices COVID-19 Vaccines Work Group: Edward Belongia, Center for Clinical Epidemiology & Population Health, Marshfield Clinic Research Institute; Dayna Bowen Matthew, George Washington University Law School; Oliver Brooks, National Medical Association; Matthew Daley, Institute for Health Research, Kaiser Permanente Colorado; Jillian Doss-Walker, Indian Health Service; Marci Drees, Society for Healthcare Epidemiology of America; Jeffrey Duchin, Infectious Diseases Society of America; Kathy Kinlaw, Center for Ethics, Emory University; Doran Fink, Food and Drug Administration; Sandra Fryhofer, American Medical Association; Jason M. Goldman, American College of Physicians; Michael Hogue, American Pharmacists Association; Denise Jamieson, American College of Obstetricians and Gynecologists; Jeffery Kelman, Centers for Medicare & Medicaid; David Kim, U.S. Department of Health and Human Services; Susan Lett, Council of State and Territorial Epidemiologists; Kendra McMillan, American Nurses Association; Kathleen Neuzil, Center for Vaccine Development and Global Health, University of Maryland School of Medicine; Sean O'Leary, American Academy of Pediatrics; Christine Oshansky, Biomedical Advanced Research and Development Authority; Stanley Perlman, Department of Microbiology and Immunology, University of Iowa; Marcus Plescia, Association of State and Territorial Health Officials; Chris Roberts, National Institutes of Health; William Schaffner, National Foundation for Infectious Diseases; Kenneth Schmader, American Geriatrics Society; Bryan Schumacher, Department of Defense; Rob Shechter, Association of Immunization Managers; Jonathan Temte, American Academy of Family Physicians; Matthew Tunis, National Advisory Committee on Immunization Secretariat, Public Health Agency of Canada; Thomas Weiser, Indian Health Service; Matt Zahn, National Association of County and City Health Officials; Rachel Zhang, Food and Drug Administration.

Corresponding author: Sara E. Oliver, yxo4@cdc.gov.

¹CDC COVID-19 Response Team; ²Epidemic Intelligence Service, CDC; ³General Dynamics Information Technology, Falls Church, Virginia; ⁴University of Arizona College of Medicine, Phoenix, Arizona; ⁵Department of Health Research Methods, Evidence, and Impact, Hamilton, Ontario; ⁶Arkansas Department of Health; ⁷Vanderbilt University School of Medicine, Nashville, Tennessee; ⁸Stanford University School of Medicine, Stanford, California; ⁹University of Washington, Seattle, Washington.

All authors have completed and submitted the International Committee of Medical Journal Editors form for disclosure of potential conflicts of interest. No potential conflicts of interest were disclosed.

Summary

What is already known about this topic?

On December 11, 2020, the Food and Drug Administration issued an Emergency Use Authorization for the Pfizer-BioNTech COVID-19 vaccine.

What is added by this report?

On December 12, 2020, after an explicit, evidence-based review of all available data, the Advisory Committee on Immunization Practices (ACIP) issued an interim recommendation for use of the Pfizer-BioNTech COVID-19 vaccine in persons aged ≥ 16 years for the prevention of COVID-19.

What are the implications for public health practice?

The recommendation for the Pfizer-BioNTech COVID-19 vaccine should be implemented in conjunction with ACIP's interim recommendation for allocating initial supplies of COVID-19 vaccines.

References

1. Food and Drug Administration. Pfizer-BioNTech COVID-19 Vaccine Emergency Use Authorization. Silver Spring, MD: US Department of Health and Human Services, Food and Drug Administration; 2020. <https://www.fda.gov/emergency-preparedness-and-response/coronavirus-disease-2019-covid-19/pfizer-biontech-covid-19-vaccine>
2. Dooling K, McClung N, Chamberland M, et al. The Advisory Committee on Immunization Practices' interim recommendation for allocating initial supplies of COVID-19 vaccine—United States, 2020. *MMWR Morb Mortal Wkly Rep* 2020;69:1857–9. PMID:33301429 <https://doi.org/10.15585/mmwr.mm6949e1>
3. Advisory Committee on Immunization Practices. ACIP meeting information. Atlanta, GA: US Department of Health and Human Services, CDC; 2020. <https://www.cdc.gov/vaccines/acip/meetings/index.html>
4. Advisory Committee on Immunization Practices. Advisory Committee on Immunization Practices (ACIP): GRADE (grading of recommendations, assessment, development and evaluation). Atlanta, GA: US Department of Health and Human Services, CDC; 2020. <https://www.cdc.gov/vaccines/acip/recs/grade>
5. Vaccines and Related Biological Products Advisory Committee. Vaccines and Related Biological Products Advisory Committee December 10, 2020, meeting: sponsor briefing document. Silver Spring, MD: US Department of Health and Human Services, Food and Drug Administration; 2020. <https://www.fda.gov/media/144246/download>
6. Vaccines and Related Biological Products Advisory Committee. Vaccines and Related Biological Products Advisory Committee December 10, 2020, meeting: FDA briefing document. Silver Spring, MD: US Department of Health and Human Services, Food and Drug Administration; 2020. <https://www.fda.gov/media/144245/download>